

# Lecture 08. Simplicial Homology Groups

Triangulation

n-Simplex  $\Delta^n = [v_0, v_1, \dots, v_n] = \left\{ (t_0, t_1, \dots, t_n) \in \mathbb{R}^{n+1} \mid t_i \geq 0, \sum_{i=0}^n t_i = 1 \right\}$ .  
 = the smallest convex subset in  $\mathbb{R}^{n+1}$  with vertices  $v_i$ .

$\Delta^n$  is ordered:  $v_0, v_1, \dots, v_n$

$$[v_1, v_0, v_2, \dots, v_n] \neq [v_0, v_1, v_2, \dots, v_n]$$

i-face of  $\Delta^n$ :  $[v_0, v_1, \dots, \hat{v}_i, \dots, v_n]$   $\partial_i$   
 $\left\{ (t_0, t_1, \dots, t_{i-1}, 0, t_{i+1}, \dots, t_n) \in \Delta^n \right\}$



$$\partial \Delta^n = \bigcup_{i=0}^n [v_0, v_1, \dots, \hat{v}_i, \dots, v_n]$$

Example.  $n=0$ ,  $\Delta^0 = [v_0]$

$$\partial \Delta^0 = \emptyset$$

$n=1$ :  $[v_0, v_1]$

$$\partial [v_0, v_1] = [v_1] - [v_0]$$

$n=2$ :  $[v_0, v_1, v_2]$

$$\begin{aligned} \partial [v_0, v_1, v_2] &= [v_1, v_2] + [v_0, v_2] + [v_0, v_1] \\ &= [v_1, v_2] - [v_2, v_0] + [v_0, v_1] \end{aligned}$$

$[v_0, v_1, v_2]$

$n=3$ :  $[v_0, v_1, v_2, v_3]$

$$\begin{aligned} \partial [v_0, v_1, v_2, v_3] &= [v_1, v_2, v_3] - [v_2, v_3, v_0] + \\ &\quad [v_0, v_1, v_3] - [v_0, v_1, v_2] \end{aligned}$$

$$[v_0, v_1, v_2, v_3]$$

In general,  $\partial [v_0, v_1, \dots, v_n] = \sum_{i=0}^n (-1)^i [v_0, v_1, \dots, \hat{v}_i, \dots, v_n]$   
formal sum

Lemma.  $\partial \partial = 0$

proof  $\partial \partial [v_0, v_1, \dots, v_n] = \partial \left( \sum_{i=0}^n (-1)^i [v_0, v_1, \dots, \hat{v}_i, \dots, v_n] \right)$

$$= \sum_{i=0}^n (-1)^i \partial [v_0, v_1, \dots, \hat{v}_i, \dots, v_n]$$

$$= \sum_{i=0}^n (-1)^i \left( \sum_{j=0}^{i-1} (-1)^j [v_0, \dots, \hat{v}_j, \dots, \hat{v}_i, \dots, v_n] + \right.$$

$$\left. \sum_{j=i+1}^n (-1)^j [v_0, \dots, \hat{v}_i, \dots, \hat{v}_j, \dots, v_n] \right)$$

$$= \sum_{0 \leq j < i \leq n} (-1)^{i+j} [v_0, \dots, \hat{v}_j, \dots, \hat{v}_i, \dots, v_n] - \sum_{0 \leq i < j \leq n} (-1)^{i+j} [v_0, \dots, \hat{v}_i, \dots, \hat{v}_j, \dots, v_n]$$

$$= 0$$


□

## Simplicial Complexes

A simplicial complex  $K$  is a finite collection of simplexes such that

- (i) If a simplex  $\sigma$  belongs to  $K$  (denoted by  $\sigma \in K$ ), then all of its faces belong to  $K$ :  $\sigma \in K \Rightarrow \sigma_i = \{v_0, \dots, \hat{v}_i, \dots, v_n\} \in K$ .
- (ii) If two simplexes  $\sigma, \tau$  satisfy  $\sigma \cap \tau \neq \emptyset$ , then  $\sigma \cap \tau$  is a common face of  $\sigma$  and  $\tau$ .

Examples.



Denote  $|K| = \bigcup_{\sigma \in K} \sigma$ , viewed as a subspace of  $\mathbb{R}^N$ ,  $N \geq 0$ .

$|K|$  is called the polyhedron associated to  $K$ .

$\dim K = \dim |K| = \max_{\sigma \in K} \{\dim \sigma\}$  is called the dimension of  $K$ .

A space  $X$  is called triangulable if there is a homeomorphism  $\varphi: |K| \rightarrow X$ .

Examples. ① polyhedra are triangulable.

② All closed/compact surfaces are triangulable.



Lemma ①  $|K| \subseteq \mathbb{R}^N$  is bounded and closed  $\Rightarrow |K|$  is compact.

② If  $|K|$  is connected, then it is path-connected.

③  $\forall x \in |K|, \exists ! \sigma \in K$ , st  $x \in \overset{\circ}{\sigma} = \text{int}(\sigma)$ . (Exercise)

## Simplicial homology groups.

Let  $K$  be a simplicial complex.

Define  $\Delta_n(K) = \mathbb{Z}\langle \text{oriented } n\text{-simplexes of } K \rangle$   
 $=$  the free abelian group generated by oriented  $n$ -simplexes  
 $= \left\{ \sum_i k_i \sigma_i \mid \sigma_i \text{ are oriented } n\text{-simplexes} \right\}$

$\partial_n: \Delta_n(K) \rightarrow \Delta_{n-1}(K)$  is a homomorphism given by

$$\partial \left( \sum_{i=0}^n (-1)^i [v_0, \dots, \hat{v}_i, \dots, v_n] \right)$$

$$\partial \left( \sum_i k_i \sigma_i \right) = \sum_i k_i \partial(\sigma_i)$$

Recall

Lemma:  $\partial_n \partial_{n+1} = 0$ :

$$0 \rightarrow \Delta_m(K) \xrightarrow{\partial_m} \Delta_{m-1}(K) \xrightarrow{\partial_{m-1}} \Delta_{m-2}(K) \xrightarrow{\partial_{m-2}} \dots \rightarrow \Delta_0(K) \rightarrow 0$$

$m = \dim K$ .

$$\partial \partial = 0 \Rightarrow \text{Im } \partial_{n+1} \subseteq \text{Ker } \partial_n$$

The quotient group  $H_n(K) := \frac{\text{Ker } \partial_n}{\text{Im } \partial_{n+1}} = \frac{Z_n(K)}{B_n(K)}$  is called

the  $n$ -th simplicial homology group of  $K$ .  $H_n(K)$  is an abelian group.

Fact:  $H_n(K)$  is a homotopy invariant of simplicial complexes:  $[a] \in H_n(K)$ ,  $[a] = a + \text{Im } \partial_{n+1}$  is called a homology class represented by  $a$ .

If  $|K| \cong |L|$ , then  $H_n(K) \cong H_n(L)$  for any  $n \geq 0$ .

$$\Delta_0(K) = \mathbb{Z} \langle \text{vertices of } K \rangle$$

$$\Delta_1(K) = \mathbb{Z} \langle \text{edges of } K \rangle$$

$$\Delta_2(K) = \mathbb{Z} \langle \text{triangles of } K \rangle$$

$$\Delta_2(K) \xrightarrow{\partial_2} \Delta_1(K) \xrightarrow{\partial_1} \Delta_0(K) \xrightarrow{\partial_0} 0$$

$$H_0(K) = \frac{\Delta_0(K)}{\partial_1(\Delta_1(K))}$$

$$\forall u, v \in \Delta_0(K), [u] = [v] \Leftrightarrow u + \text{Im } \partial_1 = v + \text{Im } \partial_1$$

$$\Leftrightarrow u - v = \partial(\sigma), \sigma \in \Delta_1(K)$$

$$\Leftrightarrow u - v = \partial([v, u] + [u, v_2] + \dots + [v_n, u])$$

$$\Leftrightarrow u \text{ and } v \text{ are two endvertices of an edge path}$$

$$\exists \gamma: [0, 1], \gamma(0) = u, \gamma(1) = v$$

$$\Leftrightarrow u \text{ and } v \text{ lie in the same (path) component.}$$

$\Rightarrow$  if  $K$  is (path) connected, then  $H_0(K) \cong \mathbb{Z}$ .

if  $K$  has  $m$  (path) components, then  $H_0(K) \cong \mathbb{Z}^m$ .

# Singular Homology

Let  $X$  be a topological space.

A map  $\sigma: \Delta^n \rightarrow X$  is called a singular  $n$ -simplex.

$$S_n(X) := \mathbb{Z} \langle \text{singular } n\text{-simplices} \rangle = \left\{ \sum_i k_i \sigma_i \mid \sigma_i: \Delta^n \rightarrow X \right\}$$

(the group of singular  $n$ -chains).

$$S_{n+1}(X) \xrightarrow{\partial_{n+1}} S_n(X) \xrightarrow{\partial_n} S_{n-1}(X)$$

$$T = [v_0, \dots, v_n]: \Delta^n \rightarrow X$$

$$\partial_n T = \sum_{i=0}^n (-1)^i T| [v_0, \dots, \hat{v}_i, \dots, v_n]$$

$$T| [v_0, \dots, \hat{v}_i, \dots, v_n]: \Delta^{n-1} = \Delta_i^n = [v_0, \dots, \hat{v}_i, \dots, v_n] \hookrightarrow \Delta^n \xrightarrow{T} X$$

Lemma:  $\partial_n \partial_{n+1} = 0$  (Exercise)

$$H_n(X) := \frac{\text{Ker } \partial_n}{\text{Im } \partial_{n+1}} = \frac{\underbrace{Z_n(X)}_{n\text{-cycle}}}{\underbrace{B_n(X)}_{n\text{-boundary}}}$$

is called the  $n$ -th singular homology group of  $X$ .

$Z_n(X)$ : elements of  $\text{Ker } \partial_n$  are called  $n$ -cycles

$B_n(X) := \text{Im } \partial_{n+1} \dots \dots$   $n$ -boundaries.

$[\alpha] = \alpha + \text{Im } \partial_{n+1} \in H_n(X)$  is called a homology class represented by a  $n$ -cycle  $\alpha$ .

Example:  $X = \{x_0\}$

$$S_n(X) = \langle \sigma: \Delta^n \rightarrow \{x_0\} \rangle = \mathbb{Z} e_n$$

$$S_n(X) \xrightarrow{\partial_n} S_{n-1}(X)$$

$$\partial e_n = \sum_{i=0}^n (-1)^i \frac{e_n| \Delta_i^n}{e_n} = \left( \sum_{i=0}^n (-1)^i \right) \cdot e_{n-1} = \begin{cases} 0 & n \text{ is odd} \\ e_{n-1} & n \text{ is even.} \end{cases}$$

$$\begin{array}{ccccccc} \mapsto S_n(X) & \xrightarrow{\partial_n} & S_{n-1}(X) & \rightarrow \dots & \rightarrow & S_2(X) & \xrightarrow{\partial_2} & S_1(X) & \xrightarrow{\partial_1} & S_0(X) & \rightarrow & 0 \\ & & & & & \text{\scriptsize } S_1 & & \text{\scriptsize } S_1 & & \text{\scriptsize } S_1 & & \\ & & & & & \xrightarrow{0} \mathbb{Z} & \xrightarrow{\cong} & \mathbb{Z} & \xrightarrow{0} & \mathbb{Z} & \rightarrow & 0 \end{array}$$

Thus  $H_0(X) \cong \mathbb{Z}$ ,  $H_i(X) \cong \mathbb{Z}/2 = 0, \forall i \geq 1$ .

Example. If  $X = \coprod_{\alpha} X_{\alpha}$ , then  $S_n(X) = S_n(\coprod_{\alpha} X_{\alpha}) \cong \bigoplus_{\alpha} S_n(X_{\alpha})$ .

$$\Delta^n \rightarrow X = \coprod_{\alpha} X_{\alpha}$$

$$S_n(X) \xrightarrow{\partial_n} S_{n-1}(X), \quad \partial_n = \bigoplus_{\alpha} \partial_{n,\alpha}$$

$$\parallel \qquad \qquad \qquad \parallel$$

$$\bigoplus_{\alpha} S_n(X_{\alpha}) \xrightarrow{\bigoplus_{\alpha} \partial_{n,\alpha}} \bigoplus_{\alpha} S_{n-1}(X_{\alpha})$$

$$\therefore H_n(X) \cong \frac{\text{Ker } \partial_n}{I_{n, \partial_{n+1}}} \cong \frac{\bigoplus_{\alpha} \text{Ker } \partial_{n,\alpha}}{\bigoplus_{\alpha} I_{n, \partial_{n+1}, \alpha}} \cong \bigoplus_{\alpha} \frac{\text{Ker } \partial_{n,\alpha}}{I_{n, \partial_{n+1}, \alpha}} = \bigoplus_{\alpha} H_n(X_{\alpha}).$$

prop.

If  $X$  is an nonempty path-connected space, then

$$H_0(X) \cong \mathbb{Z} \iff \widetilde{H}_0(X) = 0.$$

If  $X = \coprod_{i=1}^m X_i$ ,  $X_i$  are path components, then  $H_0(X) \cong \mathbb{Z}^m$

$$\text{proof.} \quad \rightarrow S_1(X) \xrightarrow{\partial_1} S_0(X) \xrightarrow{\varepsilon} \mathbb{Z} \quad \widetilde{H}_0(X) \cong \mathbb{Z}^m$$

By definition,  $H_0(X) = \frac{S_0(X)}{I_{0, \partial_1}}$

Define  $S_0(X) \xrightarrow{\varepsilon} \mathbb{Z}$ ,  $\varepsilon(\sum_i k_i \sigma_i) = \sum_i k_i$ .  $\varepsilon(\sigma) = 1$  for  $\sigma: \Delta^0 \rightarrow X$ .


- $\varepsilon$  is a group homomorphism.
- $\varepsilon$  is surjective.  $\forall k \in \mathbb{Z}$ , take  $\sigma: \Delta^0 \rightarrow X$ ,  $\varepsilon(k\sigma) = k$ .

Claim:  $I_{0, \partial_1} = \text{Ker } \varepsilon$  if  $X$  is path-connected.

$$I_{0, \partial_1} \subseteq \text{Ker } \varepsilon \iff \varepsilon \partial_1 = 0. \quad \varepsilon(\partial_1 \sigma) = \varepsilon(\sigma|_{[v_1, v_2]} - \sigma|_{[v_2, v_3]}) = 1 - 1 = 0.$$

$$\sigma: [v_1, v_2, v_3] \Delta^1 \rightarrow X$$

$\text{Ker } \varepsilon \subseteq I_{0, \partial_1}$ :  $\forall \sum_i k_i \sigma_i \in \text{Ker } \varepsilon$ ,  $\sum_i k_i = 0$ ,  $\underline{\sigma}_i = [v_i]: \Delta^0 \rightarrow X$ .  
 $x_i = \sigma_i(v_i)$ ,  $\forall i$ .



Take  $y$  to be a (base) point that is different from  $x_1, x_2, \dots, x_n$ .

Since  $X$  is path-connected, there exist paths  $\gamma_i$

$$\text{st } \gamma(0) = y, \gamma(1) = x_i$$

$\gamma_i: I = \Delta^1 \rightarrow X$  can be viewed as 1-simplex in  $X$


$$\partial \gamma_i = x_i - y.$$

Let  $\gamma = \sum_i k_i \gamma_i$ , then  $\partial \gamma = \partial(\sum_i k_i \gamma_i) = \sum_i k_i x_i - (\sum_i k_i) y = \sum_i k_i \sigma_i$ .

Lemma. (Universal property of quotient groups).


Given a homomorphism  $f: G \rightarrow H$  and a canonical projection  $\pi: G \rightarrow G/N$ .

If  $N \subseteq \text{Ker} f$ , then there is a unique homomorphism  $\bar{f}: G/N \rightarrow H$  st.  $\bar{f} \circ \pi = f$ .



Proof. Define  $\bar{f}(\pi(x)) = f(x)$ .

$N \subseteq \text{Ker} f \Rightarrow \bar{f}$  is well-defined.  $\square$



$\tilde{H}_0(X) = \text{Ker} \tilde{E} = \text{Ker} E / I_{nd_0}$  ( $= 0$  if  $X$  is path-connected)

is called the reduced 0-th singular homology group of  $X$ .

$$H_0(X) / H_0(X) \cong \mathbb{Z} \Leftrightarrow \begin{cases} H_0(X) \cong \tilde{H}_0(X) \oplus \mathbb{Z} \\ H_0(X) = 0 \text{ if } X \text{ is path-connected.} \end{cases}$$

Lemma 2. If  $G/N \cong \mathbb{Z}$ ,  $G$  and  $N$  are abelian groups, then  $G \cong N \oplus \mathbb{Z}$ .

Proof.  $0 \rightarrow N \xrightarrow{i} G \xrightarrow[\pi]{s} \mathbb{Z} \rightarrow 0$

$\mathbb{Z} = \langle 1 \rangle$ , for  $g_0 \in \pi^{-1}(1)$ , define  $s(1) = g_0 \Rightarrow \pi s = \text{id}_{\mathbb{Z}}$ .

Define  $N \oplus \mathbb{Z} \xrightarrow{(i, s)} G$

$(i, s)(x, y) = i(x) + s(y)$ .

$(i, s)$  is injective: If  $i(x) + s(y) = 0$ , then

$0 = \pi \circ i(x) + \pi \circ s(y) = 0 + y \Rightarrow y = 0$

$\therefore i(x) = 0$ . Since  $i$  is injective, we have  $x = 0$ .

$(i, s)$  is surjective:  $\forall g \in G$ ,  $s \circ \pi(g) \in G$ .

$g - s \circ \pi(g) \in \text{Ker}(\pi)$ :  $\pi(g - s \circ \pi(g)) = 0$

$x = g - s \circ \pi(g) \in N$

$y = \pi(g)$ . Then  $(i, s)(x, y) = g$ .  $\square$

$$S_n(X) \xrightarrow{\partial_0} S_{n-1}(X) \xrightarrow{\partial_1} \dots \xrightarrow{\partial_{n-1}} S_1(X) \xrightarrow{\partial_n} S_0(X) \xrightarrow{\partial_0} \mathbb{Z}$$

reduced homology groups  $\tilde{H}_n(X) =$  the homology groups of  $\{S_n(X), \partial_n; S_0(X) = \mathbb{Z}\}$   
 $\partial_0 = \varepsilon$

$$\begin{cases} \tilde{H}_n(X) \cong H_n(X) & \text{for } n > 0. \\ \tilde{H}_0(X) \cong H_0(X)/\mathbb{Z}. \end{cases}$$

• Induced homomorphisms

Let  $f: X \rightarrow Y$  be map.  $\Delta^n \xrightarrow{\sigma} X \xrightarrow{f} Y$

Define  $f_{\#}: S_n(X) \rightarrow S_n(Y)$ ,  $f(\sigma) = f \circ \sigma$ .

Lemma:  $\partial f_{\#} = f_{\#} \partial$ :

$$\begin{array}{ccc} S_n(X) & \xrightarrow{f_{\#}} & S_n(Y) \\ \downarrow \partial & & \downarrow \partial \\ S_{n-1}(X) & \xrightarrow{f_{\#}} & S_{n-1}(Y) \end{array}$$

proof  $(\partial f_{\#})(\sigma) = \partial(f \circ \sigma) = \sum_{i=0}^n (-1)^i (f \circ \sigma) | [v_0, \dots, \hat{v}_i, \dots, v_n]$   
 $= \sum_{i=0}^n (-1)^i f(\sigma | [v_0, \dots, \hat{v}_i, \dots, v_n])$   
 $= f \left( \sum_{i=0}^n (-1)^i \sigma | [v_0, \dots, \hat{v}_i, \dots, v_n] \right)$   
 $= f_{\#}(\partial \sigma) = (f_{\#} \circ \partial)(\sigma)$ .  $\square$

•  $f_{\#}: \text{Ker } \partial_n^X \rightarrow \text{Ker } \partial_n^Y$ ,  $\partial \sigma = 0 \Rightarrow \partial(f_{\#} \sigma) = f_{\#}(\partial \sigma) = 0$

$\text{Im } \partial_{n+1}^X \rightarrow \text{Im } \partial_{n+1}^Y$ ,  $f_{\#}(\partial \sigma) = \partial(f_{\#} \sigma)$

Thus  $f: X \rightarrow Y$  induces a homomorphism  $f_{\#}: H_n(X) \rightarrow H_n(Y)$ ,  $\forall n \geq 0$ .

Moreover,  $(g \circ f)_{\#} = g_{\#} \circ f_{\#}: S_n(X) \rightarrow S_n(Y) \rightarrow S_n(\mathbb{Z})$

$(\text{id}_X)_{\#} = \text{id}_{S_n(X)} = S_n(X) \rightarrow S_n(X)$

$\Rightarrow (g \circ f)_{\#} = g_{\#} \circ f_{\#}$ ,  $(\text{id}_X)_{\#} = \text{id}_{H_n(X)}$ .

$\Rightarrow$  If  $X$  is contractible, then  $\tilde{H}_n(X) = 0$ ,  $\forall n \geq 0$ .

②  $f \cong g: X \rightarrow Y$ ,  $f \circ \sigma = g \circ \sigma \Rightarrow f_{\#} = g_{\#} \Rightarrow f_{\#} = g_{\#}: H_n(X) \rightarrow H_n(Y)$ ,  $\forall n$

③ If  $X \cong Y$ , then  $H_n(X) \cong H_n(Y)$ . ( $H_n(X)$  is a top invariant.)  $\square$